

# **Structure and Sedimentology of the Dundas Group, western Tasmania.**

by

**David Selley**

B.Sc. (Hons) (University of Adelaide)




Submitted in fulfilment of the requirements for the degree of

**Doctor of Philosophy**

University of Tasmania  
Hobart

April, 1997

*This thesis contains no material which has been accepted for the award of any other degree or diploma in any tertiary institution and, to the best of my knowledge and belief, contains no copy or paraphrase of material previously published or written by another person, except when due reference is made in the text of the thesis.*

A handwritten signature in black ink, consisting of a large, stylized capital 'D' followed by a series of loops and a long, sweeping underline that extends to the left.

**David Selley**

## ABSTRACT

Middle and Late Cambrian sediments and minor intrusives of the Dundas Group form the western most exposures in the Mt Read volcanic belt of western Tasmania. The Dundas Group comprise at least 2500 m of clastic sediments with rare pumice-bearing quartz and feldspar-phyric volcanoclastics, minor quartz-feldspar porphyritic intrusives and andesitic to dacitic lavas. Volcanogenic deposits become less abundant towards the western limits of the unit, where lithologies are characterised by thick polymict conglomerate packages which occur intercalated with mudstone and thinly bedded sandstone. Structural and stratigraphic complexities of the Dundas area, with particular emphasis on the Dundas Group, provide the focus of this study.

The provenance of the Dundas Group is linked to the presently exposed basement lithologies and coeval volcanics. These include the tholeiite dominated Crimson Creek Formation, boninites, gabbros and peridotites of the Serpentine Hill Complex, older mature meta-sandstones, and the volcanics of the Mount Read volcanic 'arc'. All these sources were detected in the conglomerates and sandstones of the Dundas Group. Suite 1 samples are dominated by Crimson Creek Formation with some contamination from mature sedimentary rocks and the mafic/ultramafic complexes. Suite 1B samples have been identified as Crimson Creek Formation rather than Dundas Group on the basis of their provenance. Suite 2 samples have been derived from the mafic/ultramafic complex and a felsic source. Suite 3 has a broad range of compositions consistent with thorough mixing of detritus derived from two or more sources. Suite 4 samples are mature sandstones which have a metasedimentary source. The distinction of Dundas Group suite 4 samples from the basement sandstones is very difficult with presence of distinctive traces of chromite very useful in recognising the younger Dundas Group sandstones. Suites 1B and 4 sandstones from the eastern zone have been tentatively identified as basement lithologies faulted up within the footwall of the Rosebery Fault.

Deposition of suite 3 siliciclastics occurred during the middle Middle Cambrian in the western area, but continued into the earliest Late Cambrian in the central area. During the latter period, suite 1A sediments were being deposited along the western margin of the basin, whereas coeval finer-grained suite 3 sediments occupied central or eastern portions of the basin. This asymmetry in lithofacies distribution from west to east reflects proximal derivation of suite 1A conglomerates and sandstones from rapidly uplifted basement sources to the west, with supply of thoroughly mixed sediment to the east.

The western-most exposures of the Dundas Group involve petrographically and chemically similar conglomerate-greywacke-mudstone successions which crop out at Dundas and at Que River to the north. These range in age from latest Middle Cambrian to probable Late Cambrian and represent a marked change in basin geometry characterised by rapid basement uplift. This phase of tectonism and associated sedimentation is coeval with the onset of thrusting and molasse-type Owen Conglomerate deposition further to the

east. In the western parts of the Mt Read volcanic belt, however, basin subsidence and quiescent marine sedimentation persisted until the middle late Cambrian. This asymmetry in facies architecture and basin evolution across the Mt Read volcanic belt corresponds to diachronous Late Cambrian E-W and downwarping of thin continental crust to the west of an advancing fold and thrust belt.

The earliest deformation recognised in the Dundas Group (D<sub>1</sub>) is characterised by pre-lithification deformation features which include coherent slides along bedding parallel surfaces, chaotic zones of liquefaction, brittle fault zones and slump folds. These structures relate to syn-sedimentary seismic shock and/or gravitational collapse following basin-floor tilting.

Regional cleavage development and upright, open to tight folding correspond to the earliest phase of Middle Devonian orogenesis (D<sub>2</sub>). The S<sub>2</sub> cleavage is the dominant penetrative fabric developed throughout the Dundas region. Mesoscopic and macroscopic folds related to S<sub>2</sub> are upright to moderately inclined and shallowly to moderately plunging, however the trend of hinge lines is quite variable, ranging from NNW-SSE to NE-SW. The S<sub>2</sub> cleavage is almost always non-axial planar. The cleavage transection is interpreted to be the result of imposition of an ENE-WSW directed D<sub>2</sub> shortening axis on an earlier, pre-D<sub>2</sub> generation of NNE- to NE-trending mesoscopic and macroscopic folds. The pre-D<sub>2</sub> folds generation is tentatively correlated with a regionally developed phase of Late Cambrian E-W to NE-SW compression.

Structural relationships in the Dundas region are most complicated towards the east and culminate in tightly folded and disrupted strata positioned within the footwall of the Rosebery Fault. These rocks are characterised by the dissection of a N-S trending upright folds by an anastomosing array of steeply dipping shear zones. Where shear zones were developed in originally well stratified lithotypes with marked competency contrast, melange-type textures have formed. Domains of melange-type deformation are well exposed in the Ring River. The dominant texture common to all disrupted units is partially to completely fragmented sedimentary layers enclosed within a fine-grained and frequently fissile argillaceous matrix. Partitioning of strain and variation of deformation styles throughout the disrupted domains occurs primarily as a function of the original sedimentary lithotypes. Three broad mesoscopic styles are defined: i) "high strain" phacoid zones, ii) domains of large-scale boudinage and pinch-and-swell structure and iii) chaotic block-in-matrix structure. Bulk flattening-type strains in the Ring River melange is indicated by chocolate tablet boudinage structure and development of orthogonal extensional vein generations. The favoured interpretation for the development of the Ring River melange is the tightening and rotation of upright, shallowly plunging, NNE- to NE-trending pre-D<sub>2</sub> folds during oblique imposition of NNW to N striking D<sub>2</sub> slaty and spaced cleavages.

The melange textures formed under conditions of intense flattening strains, low effective confining pressure, marked competency contrast and relaxation of along strike boundary constraints to allow stretching in two directions. These conditions are common in accretionary environments but are not restricted to them. Similar conditions can occur in other upper crustal locations and produce melange from well-consolidated ancient rocks. The spatial relationships in the Ring River and throughout the eastern part of the Dundas region provide no support for the model that this melange was developed in an accretionary environment as suggested by Corbett and Lees (1987).

Four principal contentious issues are addressed here. Firstly the nature of the lower contact of the Dundas Group in the Ring River. This locality has been singled out as a critical test for one of the favoured tectonic models for western Tasmania. The work provided here indicates that this is a normal nonconformable contact as required by this model. The second major issue is the distribution of basement lithologies faulted into the eastern part of the map area. Major advances in the recognition of these blocks was produced by the detailed provenance studies. Thirdly, historical difficulty in constructing a coherent vertical and lateral stratigraphy in the Dundas region has been attributed to deposition within a series of sub-basins, which were periodically isolated from one another and derived material from different sediment dispersal systems. Finally the nature and significance of the melanges and disrupted zones along the Rosebery tectonic zone was given very detailed attention and the model that they are sufficient evidence to suggest a suture zone has been disproved.

## Acknowledgments

The work presented here would not have been possible without the support and guidance of my supervisor Dr Ron Berry. Throughout the final year of this project, the amount of time and energy Ron has dedicated to me will probably never be repaid. I give you my most warmest thanks. Thanks also go to the remaining academic staff of the Geology Dept., University of Tasmania for stimulating discussions and interest in my work. Particular acknowledgment must go to Drs Paul Kitto, Stuart Bull, Tony Crawford, Michael Roach and Richard Keele who have helped me significantly at various times during the course of this study.

To my dear friends from C461\* - who I miss very much, thanks for helping make this a very pleasant period of my life.

Interest in my work from geologists at the Tasmanian Department of Mines has been greatly appreciated and in particular I would like to thank Drs Tony Brown and Keith Corbett who have provided me with their time and expertise on western Tasmanian geology.

Expert technical assistance from Simon Stevens, Nathan Duhig, June Pongratz, Nilar Hlaing, and Phil Robinson as well as the administrative brilliance of Christine Higgins, Peter Cornish and Jeanette Harris has made working here very pleasant.

Special thanks go to all my fellow students for their friendship support and encouragement over the years.

Finally, to my family: Corin, my parents, my brother and sister, whose love, understanding and faith has been absolute, I am always indebted.

# TABLE OF CONTENTS

Abstract	i
Acknowledgments	iv
Table of Contents	v
List of Figures	x
List of Tables	xiii

## Chapter 1 Introduction and Regional Geology

1.1	Aims and scope of study	1
1.2	Access and Exposure	1
1.3	Regional Geology	2
1.3.1	Late Proterozoic elements ("basement")	2
1.3.2	Dundas trough Successions	3
1.3.2.1	Late Proterozoic - early Cambrian passive margin succession	3
1.3.2.2	Late early Cambrian - early Middle Cambrian mafic-ultramafic complexes	4
1.3.2.3	Middle and Late Cambrian sequences from the western Mt Read Volcanic belt	4
1.3.2.4	Late Cambrian - Early Ordovician molasse-type sequences	7
1.3.3	Lower to Middle Palaeozoic tectonic development of the dundas trough	7
1.3.3.1	Cambrian orogenesis	7
1.3.3.2	Devonian Orogenesis	10

## Chapter 2 Local Geology and Structural Relationships in Dundas Group Strata

2.1	Introduction	13
2.2	Pre-Dundas Group successions	13
2.2.1	Concert Schist and correlates of the Oonah Formation	13
2.2.2	Correlates of the Crimson Creek Formation	14
2.2.3	Mafic-Ultramafic Complexes	16

<b>2.3</b>	<b>Relationships between the mafic-ultramafic complexes and the Dundas Group</b>	<b>17</b>
<b>2.4</b>	<b>Dundas Group and correlates</b>	<b>19</b>
2.4.1	Type Area	20
2.4.2	Black Hill area	25
2.4.3	Central Area	26
2.4.4	Eastern Area	27
<b>2.5</b>	<b>Structural relationships of the Dundas Group</b>	<b>27</b>
2.5.1	Mesoscopic D <sub>2</sub> structures	28
2.5.1.1	<i>S<sub>2</sub> cleavage</i>	28
2.5.1.2	<i>Mesoscopic Folds</i>	29
2.5.2	Macroscopic structure: analysis by subarea	35
2.5.2.1	<i>Dundas township domain</i>	36
2.5.2.2	<i>Black Hill Domain</i>	38
2.5.2.3	<i>Northeast Dundas domain</i>	39
2.5.3	Summary of D <sub>2</sub> and implications for early (pre-D <sub>2</sub> ) deformation	42

## **Chapter 3 Sedimentary facies of the Dundas Group**

<b>3.1</b>	<b>Introduction</b>	<b>44</b>
<b>3.2</b>	<b>Facies 1: Conglomerates</b>	<b>44</b>
3.2.1	Subfacies 1A: Clast-supported conglomerates	45
3.2.2	Subfacies 1B: Disorganised matrix-supported conglomerate	47
3.2.3	Subfacies 1C: Open- to closed-framework mafic/ultramafic clast breccia	50
<b>3.3</b>	<b>Facies 2: Sandstones</b>	<b>53</b>
3.3.1	Subfacies 2A: Thick-bedded poorly organised sandstone	53
3.3.2	Subfacies 2B: Parallel stratified sandstones	54
<b>3.4</b>	<b>Facies 3: Sandstone-mudstone couplets</b>	<b>56</b>
3.4.1	Subfacies 3A: normally graded sandstone-mudstone couplets	56
3.4.2	Subfacies 3B: Ripple-laminated sandstone-mudstone couplets	57
<b>3.5</b>	<b>Facies 4: Fine-grained sediments</b>	<b>60</b>
<b>3.6</b>	<b>Facies 5: Immature volcanoclastic deposits</b>	<b>61</b>
3.6.1	Subfacies 5A: Feldspathic crystal-rich volcanoclastic sandstone	61



3.6.2	Subfacies 5B: Felsic vitriclastic sandstone	64
3.6.3	Subfacies 5C: Quartz-feldspathic crystal-rich sandstone	67
3.6.4	Subfacies 5D: Basalt fragment-bearing mud-matrix breccia	70
3.7	<b>Geochemical affinities of volcanoclastic deposits</b>	76

## **Chapter 4 Provenance of siliciclastics**

4.1	<b>Introduction</b>	80
4.2	<b>Chemical data sources</b>	81
4.3	<b>Sample preparation and analytical techniques</b>	82
4.4	<b>Compositional variation and classification</b>	82
4.5	<b>Suite 1</b>	83
4.5.1	Suite 1A	83
4.5.1.1	Sandstone petrography	85
4.5.1.2	Chemical composition and source characteristics	88
4.5.2	Suite 1B	89
4.5.2.1	Sandstone petrography	90
4.5.2.2	Chemical composition and source characteristics	91
4.5.3	Suite 1C	92
4.5.3.1	Sandstone Petrography	93
4.5.3.2	Chemical composition and source characteristics	94
4.5.4	Origin of the basaltic signature	96
4.5.4.1	Discrimination using MORB-normalised multi-element patterns	97
4.5.4.2	Composition and significance of detrital Cr-spinels in suite 1 sandstones	98
4.6	<b>Suite 2</b>	100
4.6.1	Distribution of samples	100
4.6.2	Sandstone petrography	100
4.6.3	Chemical composition and source characteristic	103
4.6.4	Discussion of likely sources	103
4.7	<b>Suite 3</b>	104
4.7.1	Sample distribution	105
4.7.2	Petrography of coarse-grained lithotypes	106

4.7.3	Chemical composition and source characteristics	109
4.7.4	Discussion of likely sources	110
<b>4.8</b>	<b>Suite 4</b>	<b>113</b>
4.8.1	Sample distribution	113
4.8.2	Sandstone petrography	113
4.8.3	Chemical composition and source characteristics	114
<b>4.9</b>	<b>Summary of provenance characteristics</b>	<b>116</b>
<b>4.10</b>	<b>Conclusion</b>	<b>118</b>

## **Chapter 5 Melange-Type Domains**

<b>5.1</b>	<b>Introduction and previous studies</b>	<b>120</b>
<b>5.2</b>	<b>Ring River disrupted zone</b>	<b>122</b>
<b>5.3</b>	<b>Penetrative planar fabrics associated structural elements in melange-type domains</b>	<b>123</b>
	Shear bands and shear fractures	126
	Folds	127
<b>5.4</b>	<b>Mesosopic deformation styles and geometry of inclusions in the Ring River melange</b>	<b>128</b>
5.4.1	"High strain" phacoid zones	128
5.4.2	Zones of large-scale boudinage and pinch-and-swell structure	132
	Internally massive pinch-and-swell structure and lenticular inclusions	133
	Inclusions with internal bedding laminations	136
	"Brittle" vs "ductile" inclusions: causes of variation in cross-sectional geometry	139
5.4.3	Chaotic block-in-matrix structure	140
<b>5.5</b>	<b>Microstructural fabrics and grain-scale deformation processes</b>	<b>141</b>
5.5.1	Low strain sandstone	141
5.5.2	Deformation of sandstone inclusions	142
5.5.2.1	Incipient ductile cataclastic flow	142
	Microfractures	142
	Quartz overgrowths and fibrous microstructures	143
	"Extended" quartz grains	144
	Mechanisms of grain breakage	145

5.5.2.2	Through-going brittle extensional and shear fractures	147
	Extensional fractures	147
	Shear fractures	148
	Origin of shear fractures	151
5.5.2.3	Inclusions showing transitional brittle-ductile behaviour: web structure	152
<b>5.6</b>	<b>Discussion</b>	<b>154</b>
5.6.1	Grain-scale processes	154
5.6.2	Kinematic implications of melange	157
 <b>Chapter 6 Summary and Synthesis</b>		
6.1	Introduction	160
6.2	Facies	160
6.3	Provenance	164
6.4	Structural relationships	168
6.5	Ring River Melange	170
6.6	Synthesis	173
 <b>References</b>		
		<b>177</b>
<b>Appendix A</b>	Whole Geochemical analyses of sedimentary rocks	<b>A1</b>
<b>Appendix B</b>	Electron microprobe analyses of Cr-spinels in sedimentary rocks	<b>B1</b>
<b>Appendix C</b>	Sample catalogue	<b>C1</b>